

# HEMORRHAGIC GASTROENTERITIS

## ***What is canine hemorrhagic gastroenteritis?***

Hemorrhagic gastroenteritis (HGE) is an acute disorder of dogs characterized by vomiting and bloody diarrhea, leading to depression, collapse and shock. Diarrhea is often profuse, and looks like raspberry jam. Young adult dogs of toy and miniature breeds, especially schnauzers and poodles, may be affected more frequently. Symptoms are sudden and severe.

## ***How is HGE diagnosed?***

The diagnosis of HGE is usually made based on bloodwork and symptoms. Usually, fever is not present. Some possible causes of HGE include bacterial infection or imbalance and stress. Sometimes there is no identifiable cause. Evaluation usually requires a complete blood count, biochemical analysis of the blood, urinalysis, and fecal tests. In some cases, further testing might be needed, such as x-rays, coagulation tests, or ultrasound of the abdomen.

The packed cell volume or hematocrit (measurement of red blood cells) is often above 60% with normal to low blood protein in dogs with HGE. Most normal dogs have a hematocrit of 37-55%. The elevated hematocrit provides an important clue that the patient may have HGE.

## ***What causes it?***

The exact cause of HGE remains unknown. A bacterium called *Clostridium perfringens* has been isolated from cultures of intestinal contents in some dogs with HGE, but its exact role in the syndrome has not been identified.

## ***How is it treated?***

Dogs with HGE will appear severely ill and, if untreated, may die. In most cases, the disorder appears to run its course in a few days if the dog is given appropriate supportive care. Intravenous fluid therapy and antibiotics provide the foundation of HGE therapy. Drugs for nausea are also given as needed. Food is withheld at first, and then a bland diet is introduced. Once the dog is doing well, he/she can often be switched back to a regular diet over several days. Subcutaneous fluids (given under the skin) are not usually considered adequate to meet the significant fluid requirements of most dogs with HGE, but might be considered in some mild cases.

If intravenous fluid therapy is not given, the dog's red blood count will continue to elevate due to dehydration and contraction of the spleen. In this situation, the dog is at risk for a potentially fatal clotting disorder called disseminated intravascular coagulation (DIC). Once DIC has begun, it is often irreversible and may result in death.



The good news is that if treated promptly, most patients with HGE respond very well. Recurrences are possible in an individual dog, but they are not common. Some dogs develop repeated episodes of HGE, sometimes precipitated by stressful events. Special diet, probiotics and sometimes preventative antibiotics in stressful situations can help to keep episodes to a minimum.

---

**References:**

*Blount Wendy DVM, PracticalVetMed*  
*Brooks Wendy C, DipABVP, VeterinaryPartner.com*  
*Clement Celeste ACVIM, VetCentric*  
*Homecare Handouts*  
*Ward Ernest DVM, Lifelearn Inc.*  
*Wellness Handouts*